

Bogoliubov compensation approach in QCD and in the electro-weak theory ¹

B.A. Arbuzov[†] and M. K. Volkov[‡]

[†] *Skobeltsyn Institute of Nuclear Physics of MSU, 119991 Moscow, Russia*

[‡] *Joint Institute for Nuclear Research, 141980 Dubna, Russia*

In previous works [1, 2, 3, 4, 5, 6] N.N. Bogoliubov compensation principle [7, 8] was applied to studies of spontaneous generation of effective non-local interactions in renormalizable gauge theories. In view of this one performs "add and subtract" procedure for the effective interaction with a form-factor. Then one assumes the presence of the effective interaction in the interaction Lagrangian and the same term with the opposite sign is assigned to the newly defined free Lagrangian.

Now we shall firstly demonstrate an application of the principle to the electro-weak theory. We start with EW Lagrangian with $N_{gen} = 3$ lepton and colour quark doublets with gauge group $SU(2)$. That is we restrict the gauge sector to triplet of W_μ^a only.

$$\begin{aligned}
 L = & \sum_{k=1}^3 \left(\frac{i}{2} (\bar{\psi}_k \gamma_\mu \partial_\mu \psi_k - \partial_\mu \bar{\psi}_k \gamma_\mu \psi_k) - m_k \bar{\psi}_k \psi_k + \frac{g}{2} \bar{\psi}_k \gamma_\mu \tau^a W_\mu^a \psi_k \right) + \\
 & + \sum_{k=1}^3 \left(\frac{i}{2} (\bar{q}_k \gamma_\mu \partial_\mu q_k - \partial_\mu \bar{q}_k \gamma_\mu q_k) - M_k \bar{q}_k q_k + \frac{g}{2} \bar{q}_k \gamma_\mu \tau^a W_\mu^a q_k \right) - \\
 & - \frac{1}{4} (W_{\mu\nu}^a W_{\mu\nu}^a); \quad W_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g \epsilon_{abc} W_\mu^b W_\nu^c. \quad (1)
 \end{aligned}$$

where we use the standard notations and ψ_k and q_k correspond to left leptons and quarks respectfully. We write here masses for leptons and quarks bearing in mind the ready Higgs phenomenology. In accordance to the Bogoliubov approach in application to QFT we look for a non-trivial solution of a compensation equation, which is formulated on the basis of the Bogoliubov procedure add – subtract. We have $L = L_0 + L_{int}$, where

$$L_0 = \sum_{k=1}^3 \left(\frac{i}{2} (\bar{\psi}_k \gamma_\mu \partial_\mu \psi_k - \partial_\mu \bar{\psi}_k \gamma_\mu \psi_k) - m_k \bar{\psi}_k \psi_k + \frac{i}{2} (\bar{q}_k \gamma_\mu \partial_\mu q_k - \partial_\mu \bar{q}_k \gamma_\mu q_k) - \right.$$

¹Talk at the International Bogolyubov Conference "Problems of Theoretical and Mathematical Physics", Moscow-Dubna, Russia, August 21-27 2009.

$$\partial_\mu \bar{q}_k \gamma_\mu q_k \Big) - M_k \bar{q}_k q_k \Big) - \frac{1}{4} W_{\mu\nu}^a W_{\mu\nu}^a + \frac{G}{3!} \cdot \epsilon_{abc} W_{\mu\nu}^a W_{\nu\rho}^b W_{\rho\mu}^c ; \quad (2)$$

$$L_{int} = \frac{g}{2} \sum_{k=1}^3 \left(\bar{\psi}_k \gamma_\mu \tau^a W_\mu^a \psi_k + \bar{q}_k \gamma_\mu \tau^a W_\mu^a q_k \right) - \frac{G}{3!} \cdot \epsilon_{abc} W_{\mu\nu}^a W_{\nu\rho}^b W_{\rho\mu}^c . \quad (3)$$

Here notation $\frac{G}{3!} \cdot \epsilon_{abc} W_{\mu\nu}^a W_{\nu\rho}^b W_{\rho\mu}^c$ means corresponding non-local vertex in the momentum space

$$(2\pi)^4 G \epsilon_{abc} (g_{\mu\nu} (q_\rho p_k - p_\rho q_k) + g_{\nu\rho} (k_\mu p_q - q_\mu p_k) + g_{\rho\mu} (p_\nu q_k - k_\nu p_q) + q_\mu k_\nu p_\rho - k_\mu p_\nu q_\rho) F(p, q, k) \delta(p + q + k) + \dots ; \quad (4)$$

where $F(p, q, k)$ is a form-factor and p, μ, a ; q, ν, b ; k, ρ, c are respectfully incoming momenta, Lorentz indices and weak isotopic indices of W -bosons.

Effective interaction $-\frac{G}{3!} \cdot \epsilon_{abc} W_{\mu\nu}^a W_{\nu\rho}^b W_{\rho\mu}^c$; $G = \frac{g\lambda}{M_W^2}$ is usually called anomalous three-boson interaction and it is considered for long time on phenomenological grounds.

Let us consider expression (2) as the new free Lagrangian L_0 , whereas expression (3) as the new interaction Lagrangian L_{int} . Then compensation conditions will consist in demand of full connected tree-boson vertices of the structure (4), following from Lagrangian L_0 , to be zero. This demand gives a non-linear equation for form-factor F .

Such equations according to terminology by Bogoliubov are called compensation equations. In a study of these equations it is always evident the existence of a perturbative trivial solution (in our case $G = 0$), but, in general, a non-perturbative non-trivial solution may also exist.

The goal of a study is a quest of an adequate approach, the first non-perturbative approximation of which describes the main features of the problem. Improvement of a precision of results is to be achieved by corrections to the initial first approximation.

Now in first approximation we come the following equation for $F(x)$

$$\begin{aligned} F(z) = & 1 + \frac{85 g \sqrt{N} \sqrt{z}}{96 \pi} \left(\ln z + 4 \gamma + 4 \ln 2 + \frac{1}{2} G_{15}^{31} (z_0 |_{0,0,1/2,-1,-1/2}^0) - \right. \\ & \left. - \frac{3160}{357} \right) + \frac{2}{3z} \int_0^z F(t) t dt - \frac{4}{3\sqrt{z}} \int_0^z F(t) \sqrt{t} dt - \frac{4\sqrt{z}}{3} \int_z^{z_0} F(t) \frac{dt}{\sqrt{t}} + \\ & + \frac{2z}{3} \int_z^{z_0} F(t) \frac{dt}{t} \quad z = \frac{G^2 N x^2}{1024 \pi^2} . \end{aligned} \quad (5)$$

Here $x = p^2$, $N = 2$. We introduce here an effective cut-off Y , which bounds a "low-momentum" region where our non-perturbative effects act and consider the equation at interval $[0, Y]$ under condition $F(Y) = 0$. We solve equation (5) and obtain

$$F(z) = \frac{1}{2} G_{15}^{31}(z |_{1, 1/2, 0, -1/2, -1}^0) - \frac{85 g \sqrt{N}}{512 \pi} G_{15}^{31}(z |_{1, 1/2, 1/2, -1/2, -1}^{1/2}) + C_1 G_{04}^{10}(z |_{1/2, 1, -1/2, -1}) + C_2 G_{04}^{10}(z |_{1, 1/2, -1/2, -1}). \quad (6)$$

where $G_{qp}^{mm}(z |_{b_1, \dots, b_p}^{a_1, \dots, a_q})$ is a Meijer function [9]. Constants C_1, C_2 are defined by boundary conditions. With $N = 2$ this gives

$$g(z_0) = -0.4301; \quad z_0 = 205.4254; \quad C_1 = 0.00369; \quad C_2 = 0.00582. \quad (7)$$

Note that there is also solution with large positive $g(z_0)$, which will be considered further on.

We use Schwinger-Dyson equation for W -boson polarization operator to obtain a contribution of additional effective vertex to the running EW coupling constant α_{ew} . The contribution under discussion reads after angular integrations

$$\begin{aligned} \Delta \Pi_{\mu\nu}(x) &= (g_{\mu\nu} p^2 - p_\mu p_\nu) \Pi(x); \quad x = p^2; \quad y' = q^2 + \frac{3x}{4}; \\ \Pi(x) &= -\frac{g G N}{32 \pi^2} \left(\int_{3x/4}^x \frac{F(y') dy'}{y' - x/2} \left(16 \frac{y'^3}{x^2} - 48 \frac{y'^2}{x} + 45y' - \frac{27}{2} x \right) + \right. \\ &\quad \left. + \int_x^Y \frac{F(y') dy'}{y' - x/2} \left(-3y' + \frac{5}{2} x \right) \right); \quad g = g(Y). \end{aligned} \quad (8)$$

So we have modified one-loop expression for $\alpha_{ew}(p^2)$, $x = p^2$

$$\alpha_{ew}(x) = \frac{6 \pi \alpha_{ew}(x'_0)}{6 \pi + 5 \alpha_{ew}(x'_0) \ln(x/x'_0) + 6 \pi \Pi(x)}; \quad \alpha_{ew}(x_0) = \frac{g(Y)^2}{4 \pi}; \quad (9)$$

where $x'_0 = 4/3 x_0$ means a normalization point such that $\Pi(x'_0) = 0$. Using expression (9) we calculate behaviour of $\alpha_{ew}(x)$ down to values of $x = M_W^2$ and obtain $\alpha_{ew}(M_W^2) = 0.0374$; ($\alpha_{ew}^{exp}(M_W^2) = \frac{\alpha(M_W)}{\sin_W^2} = 0.0337$). It is only 10% larger than the experimental value. We consider this result as strong confirmation of the approach.

Let us consider a contribution of effective interaction in (3) to $g - 2$ anomaly Δa . Considering contributions of the the new three-boson interaction to $W W H$ interaction we have from its conventional definition

$$G = \frac{g \lambda}{M_W^2}; \quad \lambda = -0.0151; \quad (10)$$

that agrees with experimental limitations. Then for mass of Higgs $M_H = 114 \text{ GeV}$ we obtain $\Delta a = 3.34 \cdot 10^{-9}$, that comfortably fits into error bars for well-known deviation [13] $\Delta a = (3.02 \pm 0.88) \cdot 10^{-9}$. With M_H growing Δa slowly decreases inside the error bars down to $2.67 \cdot 10^{-9}$ for $M_H = 300 \text{ GeV}$ [6].

The same procedure we apply to QCD Lagrangian with $N_f = 3$ colour quarks with gauge group $SU(3)$ and study a possibility of a spontaneous generation of effective interaction $-G_g/3! \cdot f_{abc} F_{\mu\nu}^a F_{\nu\rho}^b F_{\rho\mu}^c$, which is usually called anomalous three-gluon interaction. The second solution of Eq.(5) for $N = 3$ reads

$$z_0 = 0.01784; g(z_0) = 2.92145; C_1 = -18.8241; C_2 = 56.2171. \quad (11)$$

Then we again use Schwinger-Dyson equation for gluon polarization operator to obtain a contribution of additional effective vertex to the running strong coupling constant α_s .

So we have modified one-loop expression for $\alpha_s(p^2)$

$$\alpha_s(x) = \frac{4\pi\alpha_s(x'_0)}{4\pi + 9\alpha_s(x'_0)\ln(x/x'_0) + 4\pi\Pi(x)}; \quad x = p^2; \quad (12)$$

where x'_0 means a normalization point such that $\Pi(x'_0) = 0$. We normalize the running coupling by conditions

$$\alpha_s(x_0) = \frac{g(Y)^2}{4\pi} = 0.679185; \quad \alpha_s(x'_0) = \frac{4\alpha_s(x_0)\pi(1 + \Pi(x_0))}{4\pi + 9\alpha_s(x_0)\ln(4/3)}. \quad (13)$$

Applying the standard transformation we have

$$\alpha_s(x) = \frac{4\pi}{9 \ln \frac{x}{\Lambda^2} \left(1 + 2g\sqrt{3} \left(\alpha_s(x'_0) \ln \frac{x}{\Lambda^2} \right)^{-1} \Pi(x) \right)}; \quad \Pi(x) = 0 \text{ for } x \geq x'_0. \quad (14)$$

From here we have also $\frac{x_0}{\Lambda^2} = 1.1707002$; $G_g = \frac{6.62198}{\Lambda^2}$. Using expressions (12, 13, 14) and normalization at point M_τ , we calculate behaviour of $\alpha_s(x)$. We also use Shirkov-Solovtsov [10] method to eliminate the ghost pole, that means the following substitution in (14)

$$\alpha_s(x) = \frac{4\pi}{9} \left(\frac{1}{\ln \frac{x}{\Lambda^2}} - \frac{\Lambda^2}{x - \Lambda^2} \right) \left(1 + \frac{2g\sqrt{3}}{\alpha_s(x'_0)} \left(\frac{1}{\ln \frac{x}{\Lambda^2}} - \frac{\Lambda^2}{x - \Lambda^2} \right) \Pi(x) \right)^{-1};$$

$$\alpha_s^{(2)} \simeq 0.67 \text{ for } Q < 250 \text{ MeV}. \quad (15)$$

As the next step we apply N.N. Bogoliubov compensation principle to studies of spontaneous generation of effective non-local Nambu – Jona-Lasinio interaction. NJL model [11, 12] proves to be effective in description of low-energy hadron physics. However, the problem how to calculate parameters of the model (G_i , Λ_i ; ...) from the fundamental QCD was not solved for a long time. For the purpose we start with the conventional QCD Lagrangian with two light quarks and use again Bogoliubov procedure *add – subtract* with test interaction

$$\begin{aligned} & \frac{G_1}{2} \cdot \left(\bar{\psi} \tau^b \gamma_5 \psi \bar{\psi} \tau^b \gamma_5 \psi - \bar{\psi} \psi \bar{\psi} \psi \right) + \\ & \frac{G_2}{2} \cdot \left(\bar{\psi} \tau^b \gamma_\mu \psi \bar{\psi} \tau^b \gamma_\mu \psi + \bar{\psi} \tau^b \gamma_5 \gamma_\mu \psi \bar{\psi} \tau^b \gamma_5 \gamma_\mu \psi \right); \end{aligned} \quad (16)$$

Here notation *e.g.* $\frac{G_1}{2} \cdot \bar{\psi} \psi \bar{\psi} \psi$ means corresponding non-local vertex in the momentum space with form-factor $F_1(p_i)$

Then we again come to the corresponding compensation equation (see [3]), which has the following non-trivial solution

$$\begin{aligned} F_1(z) = & C_1 G_{06}^{40} \left(z | 1, \frac{1}{2}, \frac{1}{2}, 0, a, b \right) + C_2 G_{06}^{40} \left(z | 1, \frac{1}{2}, b, a, \frac{1}{2}, 0 \right) + \\ & C_3 G_{06}^{40} \left(z | 1, 0, b, a, \frac{1}{2}, \frac{1}{2} \right); \quad a = -\frac{1 - \sqrt{1 - 64u_0}}{4}; \quad F_1(u_0) = 1; \quad (17) \\ b = & -\frac{1 + \sqrt{1 - 64u_0}}{4}; \quad \beta = \frac{(G_1^2 + 6G_1G_2)N}{16\pi^4}; \quad z = \frac{\beta x^2}{64}; \quad u_0 = \frac{\beta m_0^4}{64}. \end{aligned}$$

Constants C_i are defined by boundary conditions. So we have the unique solution. Value of parameter u_0 and ratio of two constants G_i are also fixed $u_0 = 1.92 \cdot 10^{-8} \simeq 2 \cdot 10^{-8}$; $G_1 = \frac{6}{13} G_2$. We would draw attention to a natural appearance of small quantity u_0 .

Thus we come to effective non-local NJL interaction, which we use to obtain the description of low-energy hadron physics [2, 3, 5]. In this way we obtain expressions for all quantities under study. Analysis shows that the optimal set of low-energy parameters corresponds to $\alpha_s = 0.67$ and $m_0 = 20.3 \text{ MeV}$. We present a set of calculated parameters for these conditions including quark condensate, constituent quark mass m , parameters of light mesons

$$\begin{aligned} \alpha_s = & 0.67; \quad m_0 = 20.3 \text{ MeV}; \quad m_\pi = 135 \text{ MeV}; \quad g = 3.16; \\ m_\sigma = & 492 \text{ MeV}; \quad \Gamma_\sigma = 574 \text{ MeV}; \quad f_\pi = 93 \text{ MeV}; \\ m = & 295 \text{ MeV}; \quad \langle \bar{q} q \rangle = -(222 \text{ MeV})^3; \quad G_1^{-1} = (244 \text{ MeV})^2; \end{aligned} \quad .$$

$$\begin{aligned}
M_\rho &= 926.3 \text{ MeV}; \quad \Gamma_\rho = 159.5 \text{ MeV} \quad M_{a_1} = 1174.8 \text{ MeV}; \\
\Gamma_{a_1} &= 350 \text{ MeV}; \quad \Gamma(a_1 \rightarrow \sigma \pi)/\Gamma_{a_1} = 0.23.
\end{aligned}
\tag{18}$$

We use input quantity m_0 , while all other quantities are calculated. The overall accuracy may be estimated to be of order of 10 – 15%. The worse accuracy occurs in value M_ρ (20%).

References

- [1] *Arbuzov B.A.* Theor. Math. Phys. 2004. V.140. P.1205.
- [2] *Arbuzov B.A.* Phys. Atom. Nucl. 2006. V.69. P.1588.
- [3] *Arbuzov B.A., Volkov M.K., I.V. Zaitsev I .V.* J.Mod. Phys. A. 2006 V.21. P.5721.
- [4] *Arbuzov B.A.* Phys.Lett. B. 2007 V.656. P.67.
- [5] *Arbuzov B.A., Volkov M.K., Zaitsev I.V.* J.Mod. Phys. A. 2009. V.24. P.2415.
- [6] *Arbuzov B.A.* Eur.Phys.Journ. C. 2009. V.61. P.51.
- [7] *Bogoliubov N.N.* Soviet Phys.-Uspekhi. 1959. V.67. P. 236; Uspekhi Fiz. Nauk. 1959. V.67. P.549.
- [8] *Bogoliubov N.N.* Physica Suppl. 1960. V.26 P. 1.
- [9] *Bateman H. and Erdélyi A. Higher transcendental functions.* V. 1. New York, Toronto, London: McGraw-Hill, 1953.
- [10] *Shirkov D.V., Solovtsov I.L.* Phys. Rev. Lett. 1997, V.79. P.1209.
- [11] *Nambu Y. and Jona-Lasinio G.* Phys. Rev. 1961. V.124. P.246.
- [12] *Volkov M.K.* Fiz. Elem. Chast. Atom. Yadra 1986. V.17. P.433.
- [13] *Jegerlehner F.* Acta Phys. Polon. B. 2007. V.38. P.3021.